

The determination of α_s from lattice QCD with 2+1 flavors of dynamical quarks

C. Davies^a, A. Gray^a, M. Alford^a, E. Follana^a, J. Hein^b, P. Lepage^c, Q. Mason^c, M. Nobes^d, J. Shigemitsu^e, H. Trotter^d, M. Wingate^e, HPQCD and UKQCD collaborations.

^aDepartment of Physics and Astronomy, University of Glasgow, Glasgow G12 8QQ, UK

^bSchool of Physics, University of Edinburgh, Edinburgh EH9 3JZ, UK

^cNewman Laboratory of Nuclear Studies, Cornell University, Ithaca NY 14853, USA

^dDepartment of Physics, Burnaby, British Columbia V5A 1S6, CANADA

^eDepartment of Physics, the Ohio State University, OH 43210, USA

We describe the first lattice determination of the strong coupling constant with 3 flavors of dynamical quarks. The method follows previous analyses in using a perturbative expansion for the plaquette and Upsilon spectroscopy to set the scale. Using dynamical configurations from the MILC collaboration with 2+1 flavors of dynamical quarks we are able to avoid previous problems of having to extrapolate to 3 light flavors from 0 and 2. Our result agrees with our previous work: $\alpha_s^{\overline{MS}}(M_Z) = 0.121(3)$.

1. INTRODUCTION

Precise determinations of the strong coupling constant are important for strong interaction phenomenology and searches for new physics. Lattice QCD provides a good methodology for such a determination [1].

2. IMPROVED STAGGERED QUARKS

The staggered quark formulation allows for much faster simulation of dynamical quarks than other formulations. It maintains a remnant chiral symmetry at the cost of not entirely removing the doublers, which appear as additional ‘tastes’ of quarks. The presence of the doublers gives rise to large lattice artefacts through taste-changing interactions. The realisation that these can be systematically reduced by a Symanzik improvement programme opens the way to generating ensembles with light dynamical quarks and small lattice artefacts. The MILC collaboration have generated a number of dynamical ensembles using the Asqtad action for the dynamical quarks

and either 2 dynamical flavors, 3 degenerate dynamical flavors or 2 degenerate plus 1 heavier dynamical flavors, which should be close to the real world [2]. In the 2+1 flavor case they have been able to reduce the light dynamical mass to 1/5 of the heavier one (which is approximately at m_s).

The Asqtad staggered action includes a set of staples in place of the simple 1-link derivative, to suppress gluon exchange with momentum π/a which couples the tastes together. These staples include a correction for low-momentum a^2 artefacts introduced by this procedure. In addition there is a 3-link derivative (Naik) term to improve rotational discretisation errors through a^2 . The Asqtad action then has $\alpha_s a^2$ rotational errors and $\alpha_s^2 a^2$ taste-symmetry breaking interactions. This is combined in MILC ensembles with a gluon action improved to the level of $\alpha_s^2 a^2$ discretisation errors.

We have calculated the Υ spectrum on these ensembles using the NRQCD action for the b quark [3].

3. NRQCD

The physics of the bottomonium (Υ) system is that of non-relativistic b quarks ($v^2/c^2 \approx 10\%$) and so the NRQCD action is appropriate [4]. We use a Hamiltonian correct through $a^2 v^4/c^4$ where all terms are tadpole-improved using an estimate of the Landau gauge link [3]. For the purposes of determining the lattice spacing to set the scale for α_s we need only the radial and orbital excitation energies. These are experimentally well-determined and have the advantage that they are to a good approximation independent of heavy quark mass so that sophisticated tuning of the b mass is not necessary. In addition they are good probes of the dynamical quark content of the theory because the Υ has no valence light quarks.

The leading contribution to the radial and orbital splittings is at $\mathcal{O}(v^2/c^2)$ and so the expected systematic error from higher order (e.g. $\alpha_s v^4/c^4$ or v^6/c^6) terms not included in the NRQCD action is expected to be at the 1% level. The splittings can be determined very precisely on the lattice because of the speed of NRQCD correlator calculations.

4. DETERMINATION OF α_s

We use the perturbative expansion of the plaquette in terms of a physically well-motivated coupling, $\alpha_V(q)$, defined from the heavy quark potential:

$$V(q) = -\frac{16\pi}{3q^2} \alpha_V(q). \quad (1)$$

We then define $\alpha_P(q)$ by the following expansion of the logarithm of the plaquette, where α_P is equal to α_V to the order (second) to which we are working here.

$$-\ln(\text{plaq}) = 3.0682(2)\alpha_P(q^*)[1 + \alpha_P\{-0.770(4) - 0.09681(9)n_f\}] \quad (2)$$

The numbers in brackets indicate numerical uncertainties in the perturbative coefficients from the numerical integration involved in their calculation. This perturbative expansion is for 1-loop Symanzik-improved gluon action using tadpole-improvement with u_0 from the plaquette and n_f




Figure 1. $\alpha_P(8.2\text{GeV})$ for various MILC ensembles with differing numbers of dynamical flavors. Two different quantities are used to set the scale: the Υ 1P-1S splitting and the Υ 2S-1S splitting. Errors from setting the scale are included.

flavors of Aqtad improved staggered quarks, as appropriate to the MILC ensembles. The scale, q^* , is set using the BLM scheme and is $3.33/a$ in this case. The perturbative expansion is well-behaved and shows small corrections for non-zero n_f .

α_P can be converted to $\alpha_{\overline{MS}}$ using the equality of α_P and α_V and the continuum relation between α_V and $\alpha_{\overline{MS}}$.

$$\alpha_{\overline{MS}}^{(n_f)}(q) = \alpha_P^{(n_f)}(e^{5/6}q)[1 + 2\frac{\alpha_P}{\pi} + \mathcal{O}(\alpha_P^2)] \quad (3)$$

5. RESULTS

Figure 1 shows results for α_P evolved to a common (and arbitrarily chosen) physical scale of 8.2 GeV for various MILC ensembles. Notice the clear separation of results for $n_f = 0, 2$ and 3.

The figure shows results using either the Υ orbital splitting or radial splitting to set the lattice spacing. The orbital splitting is calculated from

the energy difference between the 1^1P_1 (h_b) and 1^3S_1 (Υ) states on the lattice. The experimental result for the 1^1P_1 mass is not known and so is taken as the spin-average of the masses of the 1^3P (χ_b) states. The difference is expected to be very small. The radial splitting is the difference in energies of the 2^3S_1 (Υ') and 1^3S_1 states, which is experimentally well-known.

The value for $\alpha_P(8.2\text{GeV})$ agrees well between the two determinations on the 2+1 dynamical ensemble but not on the others. This reflects the fact that the lattice spacing will depend on the quantity chosen to fix it if the vacuum does not have dynamical content which is close enough to the ‘real world’.

We can take the result for α_P on the 2+1 dynamical ensemble, convert to $\alpha_{\overline{MS}}$ and run to M_Z . The result for $\alpha_{\overline{MS}}^{(5)}$ is then 0.121(3). The dominant source of error (0.002) is the perturbative error from unknown higher orders in perturbation theory in eq. 3. This result agrees with the world average and with our previous results using dynamical (unimproved) staggered quarks [1].

Other sources of systematic error give small effects. We can see from Figure 1 that the $n_f = 3$ results show very little dependence on dynamical light quark mass. Extrapolating α_P with m_π^2 would change the final result by less than 0.001.

The lattice spacing dependence of the result is expected to be very small but we currently have Υ results only on the coarser of the MILC set of ensembles. We can estimate the dependence from other results, however. Figure 2 shows the dependence of α_P in the quenched approximation on a^2 for improved and unimproved glue [5]. Here we have used r_0 to set the scale (and quote α_P at a scale of $15/r_0$) because it can be fixed very precisely. Its disadvantage is that it is not clear how to convert it to a physical result. The figure shows that unimproved glue has very small dependence on a and improved glue (appropriate to the MILC ensembles) even less. We fold this lattice spacing dependence along with quark mass effects above and NRQCD systematics into a final error of 0.003.




Figure 2. α_P in the quenched approximation at a scale of $15/r_0$ for improved and unimproved glue.

6. CONCLUSIONS

The first calculation of α_s with 2+1 flavours of dynamical quarks gives results in agreement with previous work, but without extrapolating in n_f . Further improvements, including the calculation of the next order in perturbation theory, are planned to reduce the error even further.

Acknowledgements

We are grateful to the MILC collaboration for the use of their configurations. This work was supported by the DoE, the EU IHP programme, NSF, NSERC and PPARC.

REFERENCES

1. C. Davies *et al*, Phys. Rev. D**56** (1997) 2755.
2. MILC collaboration, Phys. Rev. D**64** (2001) 054506.
3. A. Gray *et al*, these Proceedings.
4. G. P. Lepage *et al*, Phys. Rev. D**46** (1992) 4052.
5. We thank Steve Gottlieb for providing the unpublished MILC results for this plot.